

UTILIZATION OF MICROWAVE SPECTROSCOPY TO IDENTIFY AND PROBE REACTION DYNAMICS OF HSNO, A CRUCIAL BIOLOGICAL SIGNALING MOLECULE

MATTHEW NAVA, *Department of Chemistry, MIT, Cambridge, MA, USA*; MARIE-ALINE MARTIN-DRUMEL, *Atomic and Molecular Physics, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA*; JOHN F. STANTON, *Department of Chemistry, The University of Texas, Austin, TX, USA*; CHRISTOPHER CUMMINS, *Department of Chemistry, MIT, Cambridge, MA, USA*; MICHAEL C MCCARTHY, *Atomic and Molecular Physics, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA*.

Thionitrous acid (HSNO), a potential key intermediate in biological signaling pathways, has been proposed to link NO and H₂S biochemistries. Its existence and stability in vivo, however, remain controversial. By means of Fourier-transform microwave spectroscopy, we establish that HSNO is spontaneously formed in high concentration when NO and H₂S gases are simply mixed at room temperature in the presence of metallic surfaces. Our measurements reveal that HSNO is formed with high efficiency by the reaction H₂S and N₂O₃ to produce HSNO and HNO₂, where N₂O₃ is a product of NO disproportionation. These studies also suggest that further reaction of HSNO with H₂S may form HNO and HSSH. The length of the S–N bond has been derived to high precision from isotopic studies, and is found to be unusually long, 1.84 Å – the longest S–N bond reported to date for an SNO compound. The present structural and reactivity investigations of this elusive molecule provide a firm foundation to better understand its physiological chemistry and propensity to undergo S–N bond homolysis in vivo.